

High prevalence of methicillin-resistant coagulase-negative staphylococci isolated from a university environment in Thailand

Rathanin Seng,¹ Udomluk Leungtonkam,¹ Rapee Thummeepak,¹
Wassana Chatdumrong,^{1,2} Sutthirat Sitthisak^{1,2*}

¹Department of Microbiology and Parasitology, Faculty of Medical Sciences, Naresuan University, Phitsanulok, Thailand.

²Centre of Excellence in Medical Biotechnology, Faculty of Medical Sciences, Naresuan University, Phitsanulok, Thailand

Received 22 March 2017 · Accepted 28 April 2017

Summary. The present study was conducted to isolate and characterize the molecular epidemiology of the methicillin-resistant staphylococci in the general university environment, where all five locations; the library, restrooms, canteens, computer rooms and outdoor surfaces, are in common use by a large population of students. We used Mannitol Salt Agar (MSA) supplemented with 4 µg/ml of oxacillin to screen the methicillin-resistant staphylococci. The species level was identified by PCR of *rdr* (*Staphylococcus epidermidis*), *groESL* (*Staphylococcus haemolyticus*) and *nuc* (*Staphylococcus aureus* and *Staphylococcus warneri*) genes and DNA sequencing of *tuf* and *dnaJ* genes. The susceptibility patterns of the isolates were determined using the disk diffusion method. Antibiotic and disinfectant resistance genes, together with SCCmec types, were detected by the PCR method. The methicillin resistant-staphylococci were isolated from 41 of 200 samples (20.5%), and all of them were found to be methicillin-resistant coagulase negative staphylococci (MR-CoNS). The library had the highest percentage of contamination, with 43.3% of the samples found to be contaminated. All isolates belonged to 6 different species including *S. haemolyticus*, *S. epidermidis*, *S. warneri*, *S. cohnii*, *S. saprophyticus* and *S. hominis*. The antimicrobial resistance rates were highest against penicillin (100%), then cefoxitin (73.1%), erythromycin (73.1%) and oxacillin (68.3%). Altogether, the isolates were approximately 61.0% multidrug resistant (MDR), with the *S. epidermidis* isolates being the most multidrug resistant ($P < 0.05$). The prevalence of the *qacA/B* gene was detected in 63.4% of the isolates, and SCCmec could be typed in 43.9% (18/41) of the isolates. The type range was: II (n = 1), IVd (n = 1), I (n = 2), V (n = 6), IVa (n = 8) and untypeable (n = 23). This result indicates that these university environments are contaminated with methicillin-resistant coagulase negative staphylococci that carry various SCCmec types and high rate of disinfectant resistance genes. [Int Microbiol 20(2):65-73 (2017)]

Keywords: *Staphylococcus* spp. · methicillin-resistant coagulase negative staphylococci · drug resistance · gene *qacA/B* · Phitsanulok (Thailand)

Introduction

Staphylococci, particularly *Staphylococcus aureus*, *S. epidermidis*, *S. saprophyticus*, *S. haemolyticus*, *S. hominis*, and

S. lugdunensis are medically important pathogens which cause nosocomial and community infections [29]. Staphylococci are classified into coagulase-positive staphylococci known as *S. aureus* and coagulase-negative staphylococci (CoNS) such as *S. epidermidis*, *S. saprophyticus*, *S. haemolyticus*, *S. hominis* and another 49 species [28]. Most strains of these bacteria have developed methicillin resistance and

* Corresponding author: Sutthirat Sitthisak
E-mail: sutthirats@nu.ac.th

are therefore called methicillin-resistant *S. aureus* (MRSA) and methicillin resistant coagulase negative staphylococci (MR-CoNS), which constitute a major health problem. More recently, the number of reports of community-acquired MRSA (CA-MRSA) has been rapidly increasing. However, infections caused by MR-CoNS in the community have not been reported, but the gene transfer from these bacteria to MRSA has been identified [23].

The resistance in MRSA and MR-CoNS is caused by the acquisition of the *mecA* gene that encodes a modified penicillin-binding protein 2a (PBP2a) which has a low binding affinity for all beta-lactam antibiotics. The *mecA* gene is located within the *mec* operon carried by staphylococcal cassette chromosome *mec* (SCC*mec*). SCC*mec* are classified into 11 different types and various subtypes [46]. SCC*mec* type I, II and III are carried by hospital-acquired MRSA (HA-MRSA) and SCC*mec* type IV, V and VI are carried by community-acquired MRSA (CA-MRSA) [6]. In contrast to MRSA, the distribution of SCC*mec* types in MR-CoNS is varied, depending on the human host and the geographical locations from where the isolates were obtained [40]. Moreover, the variety of other antibiotic and disinfectant resistance genes such as erythromycin resistance genes (*erm*) and the quaternary ammonium compound resistance gene (*qacA/B*) were either identified in plasmids of, or by mobile genetic elements of, staphylococci [44].

Recent studies have reported that a wide variety of different high-touch environmental surfaces in public facilities, universities, microbiological and computer laboratories, day-care centers, prisons and clinics are a potential reservoir of MRSA and drug resistance genes [7]. However, environmental colonization to the spread of MR-CoNS is poorly reported, although Widerström and coworkers [38] found that the hospital-acquire methicillin-resistant *S. epidermidis* was spread from environmental fomites to patients in Intensive Care Units. Xu, Mkrtchyan and Cutler [40] reported that 21% of hotel samples were contaminated with MR-CoNS, and Mkrtchyan and coworkers [20] found that staphylococci are the most predominant bacteria in non-hospital restrooms, in the UK. Moreover, 23 MR-CoNS were isolated from 7 beach sites in Washington State by Soge and coworkers [33]. We have been able to find only these three papers published on the matter, therefore there is little information available on the prevalence and molecular epidemiology of methicillin-resistant staphylococci isolated from non-hospital environments, and this is especially so in the case of Thailand.

Therefore, exploration of this information has become an important issue demanding investigation, which was the purpose of the current study. We isolated and identified methicillin-resistant staphylococci at the specie level, from a university environment in Thailand, and determined the antimicrobial susceptibility pattern, detected the antibiotic and disinfectant resistance genes, and characterized the SCC*mec* types of all isolates.

Materials and methods

Population and samples. The sample size of the study was calculated using the Sample Size Determination in Health Studies Software of the World Health Organization [17]. A total of 200 swab samples were randomly collected from 5 locations of a university community in Thailand, including computer rooms (computer mouse, computer earpieces, computer keyboards and computer power buttons) (n = 40); restrooms (door handles, washbasins, washbasin areas, urinary taps and toilets) (n = 50), the library (books, escalators and tables) (n = 30), canteens (tables, bank notes and coins used for payment, ATM machines and water dispensers) (n = 40), and outdoor surfaces (handrails, exercise machines, public buses) (n = 40). The sample collection was carried out from September to December 2015 with temperature ranging from 31 °C to 32 °C at the study area.

Bacterial isolation of methicillin-resistant staphylococci.

The samples were collected using cotton swabs soaked in 0.85% normal saline, and then placed in transfer media (2% of skim milk powder, 3% of tryptone soya broth (TSB), 0.5% glucose and 10% glycerol). The swab samples were enriched in TSB with shaking at 180 rpm at 37 °C for 18–24 h. Then, one loopful of overnight culture was streaked on MSA with 4 µg/ml of oxacillin and incubated at 30 °C for 48–72 h for a primary screening for methicillin-resistant staphylococci base on the method of Lally et al. [15]. All colonies were selected for further identification using Gram's stain, catalase and coagulase tests. All isolates were subsequently confirmed as staphylococci by the PCR method amplified by 16S rRNA specific primers [13]. Methicillin-resistance was further confirmed by oxacillin disk (1 µg), cefoxitindisk (30 µg) and the PCR method to detect *mecA* gene. *Staphylococcus aureus* COL was used as positive control of this method.

Identification of methicillin-resistant staphylococci species.

Staphylococcus aureus, *S. epidermidis*, *S. haemolyticus* and *S. warneri* were distinguished from other species by the PCR method based on the specific primer of *nuc* (*S. aureus* and *S. warneri*), *rdr* (*S. epidermidis*) and *groESL* (*S. haemolyticus*) as described by Schmidt et al. [28]. A specific gene of each species was sequenced to ensure the absence of bias in our method. The remaining isolates that could not be identified by PCR were identified by *tuf* and *dnaJ* gene sequencing using Sanger Sequencing Method, according to the methods described by Loonen et al. [16] and Shah et al. [30]. The PCR products were purified using an RBC purification kit and sequenced using Applied Biosystems. Sequence similarities of *tuf* and *dnaJ* genes > 97% were used to identify isolates at the species level. The primer sets of *nuc*, *rdr*, *groESL*, *tuf* and *dnaJ* genes are shown in Table 1.

Table 1. Primers for antibiotic resistance gene amplification and bacterial identification

Primers	Oligonucleotide sequence (5'-3')	Tm (°C)	Product size (bp)	Reference
<i>nuc-F</i> (<i>S. aureus</i>)	TCGCTTGCTATGATTCTGG	55.2	359	[27]
<i>nuc-R</i> (<i>S. aureus</i>)	GCCAATGTTCTACCATAGC	55.2		
<i>nuc-F</i> (<i>S. hominis</i>)	TACAGGGCCATTTAAAGACG	56.4	177	[10]
<i>nuc-R</i> (<i>S. hominis</i>)	GTTTCTGGTGTATCAACACC	56.4		
<i>nuc-F</i> (<i>S. warneri</i>)	CGTTTGTAGCAAAAACAGGGC	58.4	999	[10]
<i>nuc-R</i> (<i>S. warneri</i>)	GCAACGAGTAACCTTGCCAC	60.5		
<i>rdr-F</i>	AAGAGCGTGGAGAAAAGTATCAAG	61.8	130	[32]
<i>rdr-R</i>	TCGATACCATCAAAAAGTTGG	61.8		
<i>groESL-F</i>	GGTCGCCTTAGTCGGAACAAT	57.8	271	[4]
<i>groESL-R</i>	CACGAGCAATCTCATCACCT	57.8		
<i>tuf-F</i>	CCAATGCCACAAAACCTCGTGA	58.4	480	[16]
<i>tuf-R</i>	CAGCTTCAGCGTAGTCTAATAATTTACG	65.7		
<i>dnaJF</i>	GCCAAAAGAGACTATTATGA	52.3	920	[30]
<i>dnaJR</i>	ATTGYTTACCYGTTTGTGTACC	56.6		
<i>mecA-F</i>	TGGCTATCGTGTACAATCG	58	310	[26]
<i>mecA-R</i>	GTTCTCTCATAGTATGACGTCC	58		
<i>ermA-F</i>	AAGCGGTAAACCCCTCTGA	56.7	190	[34]
<i>ermA-R</i>	TTCGCAAATCCCTTCTCAAC	55.2		
<i>ermB-F</i>	AATCGTCAATTCCTGCATGT	55.9	142	[3]
<i>ermB-R</i>	TAATCGTGGAAATACGGGTTTG	55.9		
<i>ermC-F</i>	AATCGTCAATTCCTGCATGT	53.2	299	[34]
<i>ermC-R</i>	TAATCGTGGAAATACGGGTTTG	55.9		
<i>qacA/B-F</i>	GCAGAAAGTGCAGAGTTCCG	57.3	361	[44]
<i>qacA/B-R</i>	CCAGTCCAATCATGCCTG	56.1		

Determination of antimicrobial susceptibility patterns. Antimicrobial susceptibility of all isolated methicillin-resistant staphylococci was tested using a standard disk diffusion test against fifteen antibiotics: penicillin (P, 10 units), clindamycin (DA, 2 µg), chloramphenicol (C, 30 µg), gentamicin (CN, 10 µg), erythromycin (E, 15 µg), cefoxitin (FOX, 30 µg), sulfamethoxazole/trimethoprim (SXT, 1.25/23.75 µg), oxacillin (OX, 1 µg), vancomycin (VA, 30 µg), rifampicin (RD, 5 µg), linezolid (LZD, 30 µg), mupirocin (MUP, 5 µg), ciprofloxacin (CIP, 5 µg), fusidic acid (FD, 10 µg) and novobiocin (NV, 5 µg). The plates were incubated at 35 °C for 24 h. *Staphylococcus aureus* NCTC10442 was used as positive control and the results were interpreted according to the guidelines of the Clinical and Laboratory Standards Institute 2014. All isolates were categorized as MDR when they were resistant to at least three classes of antibiotics [19].

Detection of antibiotic and disinfectant resistance genes. The methicillin-resistance gene (*mecA*) was detected according to the method described by Kitti, Boonyonying and Sitthisak [12]. *Staphylococcus aureus* COL was used as positive control of this detection. Other antibiotic resistant genes, including *erm(A)*, *erm(B)*, *erm(C)* and *qacA/B*, were detected by PCR modified from the method described in [1,41,44]. The primers used are shown in Table 1. The absence of bias was ensured by the sequencing of each

gene in the representative isolates. All PCR products were visualized using gel electrophoresis with 1% agarose gel stained with 0.5% ethidium bromide.

Characterization of SCC*mec* types. SCC*mec* types of all isolates were characterized according to the method of Zhang et al. [43]. *Staphylococcus aureus* NCTC10442, *S. aureus* JCSC10442, *S. aureus* WIS and *S. aureus* isolated from our previous study [36] were used as reference strains of SCC*mec* Type I, II, IVa, IVb and V. The amplicons were visualized using gel electrophoresis with 1% agarose gel stained with 0.5% ethidium bromide.

Statistical analysis. All data was analyzed using Stata 12.0 (Stata Corporation, USA). The analysis of frequency (Chi-square test; $P < 0.05$) was used as the statistic to compare the MDR and antimicrobial susceptibility patterns among MR-CoNS species. The association between MR-CoNS prevalence and environmental locations was calculated using logistic regression ($P < 0.05$).

Results

The prevalence of methicillin-resistant staphylococci. We used MSA, supplemented with 4 µg/ml of oxa-

Table 2. The association of MR-CoNS prevalence with environmental locations

Locations ^a	N	Positive MR-CoNS (%)	Negative MR-CoNS (%)	OR	95% CI	P-value
LB	30	13 (43.3)	17 (56.7)			
RR	50	14 (28.0)	36 (72.0)	0.50	0.19, 1.31	
CN	40	8 (20.0)	32 (80.0)	0.32	0.11, 0.94	
CR	40	5 (12.5)	35 (87.5)	0.18	0.05, 0.60	
US	40	1 (2.5)	39 (97.5)	0.03	0.00, 0.27	<0.001*

^aLB: Library, RR: Restroom, CN: Canteen, CR: Computer room and US: Outdoor surfaces
OR: odds ratio. CI: confidence interval. *Significant at $P < 0.001$

cillin, to screen for methicillin-resistant staphylococci and confirmed by oxacillin disk (1 µg), cefoxitin disk (30 µg) and *mecA* gene detection. Among the 200 samples, 41 (20.5%) were identified as methicillin-resistant staphylococci, while all of them were MR-CoNS. The library was the most contaminated, with 43.3% of the locations tested showing staphylococci contamination. The next most contaminated were the restrooms (28%), canteens (20%), computer rooms (12.5%) and outdoor surfaces (2.5%).

Logistic regression was performed to analyze the association between the environmental locations and MR-CoNS

prevalence. Overall, the environmental locations tested in our study were significantly associated with colonization of MR-CoNS ($P < 0.001$). More precisely, the library was the most contaminated region (Table 2).

Species distribution. The specie level of all isolates was identified by our combined method of biochemical test, PCR and DNA sequencing. All 41 isolates of MR-CoNS belonged to 6 different species including *S. haemolyticus* (41.5%), *S. epidermidis* (36.6%), *S. warneri* (12.2%), *S. cohnii* (4.9%), *S. saprophyticus* (2.4%) and *S. hominis* (2.4%) (Fig. 1).




Fig. 1. Prevalence of MR-CoNS and species distribution by environmental locations (LB: Library, RR: Restroom, CN: Canteen, CR: Computer room, and US: Outdoor surfaces.)




Fig. 2. Differences among MDR and NON MDR pattern of each species. *Significant at $P < 0.05$.

Antimicrobial susceptibility pattern. All isolates were resistant to penicillin (100%), cefoxitin (73.1%), erythromycin (73.1%), oxacillin (68.3%), sulfamethoxazole/trimethoprim (29.3%), fusidic acid (22.0%), clindamycin (14.6%), ciprofloxacin (12.2%), chloramphenicol (9.8%), novobiocin (9.8%), gentamicin (4.9%), rifampicin (2.4%), mupirocin (2.4%). All isolates were susceptible to vancomycin and linezolid. We found most of the isolates (approximately 61.0%) were MDR. We divided all 41 isolates into three species groups, *S. haemolyticus*, *S. epidermidis* and other species. Then, we compared the resistance rate of each antibiotic among these species using chi-square test. Significantly, the prevalence of MDR in each group was different ($P < 0.05$). *Staphylococcus epidermidis* and other species (*S. warneri*, *S. cohnii*, *S. hominis* and *S. saprophyticus*) were more associated with MDR than was *S. haemolyticus* ($P < 0.05$) (Fig. 2). *Staphylococcus haemolyticus* was more resistant to clindamycin than were *S. epidermidis* and other species ($P < 0.05$) (Table 3).

Antibiotic and disinfectant resistance genes.

Gene *mecA* was detected in all MR-CoNS isolates by the PCR method. Due to the high resistance to erythromycin, its resistance genes: *erm(A)*, *erm(B)* and *erm(C)* were also detected. Only *erm(C)* was detected in 14.6% of isolates and 20.0% of

erythromycin resistant isolates, indicating the low correlation between the phenotypic pattern and the causative genes. We also detected *qacA/B* gene, the gene encoding resistance to several antiseptics and disinfectants. A high proportion (63.4%) of all isolates was positive with this gene.

SCCmec types. All isolates were subjected to characterization by their SCCmec types using multiplex PCR. Among all the isolates, 43.9% (18/41) were characterized as SCCmec type I (n = 2), II (n = 1), IVa (n = 8), IVd (n = 1) and V (n = 6). We found 56.1% (23/41) were untypeable SCCmec types. The distribution of SCCmec types in each species is shown in Table 4.

Discussion

To strengthen the understanding about the dissemination of methicillin-resistant staphylococci within non-health care environments, we examined isolates from a university environment in Thailand, and identified the specie levels in those isolates. Most of the MR-CoNS (43.3%) isolates in the study were obtained from items in the library, such as books and study tables. Hence, MR-CoNS can be easily spread by the simple act of reading a book at a table in the library.

Table 3. The comparison of *qacA/B* gene and antimicrobial resistance patterns in each MR-CoNS species

Antibiotics	<i>S. haemolyticus</i> n = 17 (%)	<i>S. epidermidis</i> n = 15 (%)	Other species n = 9 (%)	Total n = 41 (%)	P-value
<i>qacA/B</i>	12 (70.6)	9 (60.0)	5 (55.6)	26 (63.4)	0.767
Oxacillin	12 (70.6)	12 (80.0)	4 (44.4)	28 (68.3)	0.187
Cefoxitin	12 (70.6)	13 (86.7)	5 (55.6)	30 (73.1)	0.238
Erythromycin	13 (76.5)	10 (66.7)	7 (77.8)	30 (73.1)	0.773
Sulfamethoxazole/ Trimethoprim	4 (23.5)	3 (20.0)	2 (22.2)	9 (22.0)	0.971
Chloramphenicol	0 (0.0)	2 (13.3)	2 (22.2)	4 (9.6)	0.162
Rifampicin	0(0.0)	1 (6.7)	0 (0.0)	1 (2.4)	0.411
Gentamicin	1 (5.9)	1 (6.7)	0 (0.0)	2 (4.9)	0.740
Fusidic acid	5 (29.4)	1 (6.7)	0(0.0)	6 (14.6)	0.071
Clindamycin	10 (58.8)	1 (6.7)	1 (11.1)	12 (29.3)	<0.05*
Mupirocin	0 (0.0)	0 (0.0)	1 (11.1)	1 1 (2.4)	0.162
Novobiocin	1 (5.9)	3 (20.0)	0 (0.0)	4 (9.6)	0.218
Ciprofloxacin	1(5.9)	2 (13.3)	2 (22.2)	5 (12.2)	0.473
Vancomycin	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	NA
Linezolid	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	NA

*Significant at $P < 0.05$.

NA = not analyzed.

Using our screening method, none of the isolates was identified as MRSA. Like in a previous report, on a study conducted in India, MRSA was not recovered from a hospital environment [31]. To ascertain the specie distribution of MR-CoNS isolates in this study, we identified that 6 different species were presented in all of the isolates (Fig. 1). These species are similar to the CoNS obtained from non-

hospital environments [20,40], clinical specimens [14,25], healthy adult volunteers [2] and chicken meat [22]. However, the prevalence of each species was different among these specimens. *Staphylococcus epidermidis* was categorized as the predominant species of CoNS isolated from clinical and commensal samples, while *S. haemolyticus* was found to have the highest prevalence in non-healthcare environments.

Table 4. Distribution of SCCmec types among MR-CoNS isolated from the university environment. The values in the table indicate the number of each SCCmec type in each species

MR-CoNS Species	SCCmec					
	I	II	IVa	IVd	V	Untypeable
<i>S. haemolyticus</i>	0	1	1	0	3	12
<i>S. epidermidis</i>	0	0	4	1	1	9
<i>S. warneri</i>	0	0	3	0	2	0
<i>S. cohnii</i>	2	0	0	0	0	0
<i>S. saprophyticus</i>	0	0	0	0	0	1
<i>S. hominis</i>	0	0	0	0	0	1
Total	2	1	8	1	6	23

In contrast, the species distribution on our study was different from CoNS contaminating fermented food and starter cultures [42].

More than 60% of all isolates were resistant to penicillin, cefoxitin, and erythromycin. Two notable antimicrobial resistance patterns of these isolates were found. First, 61.0% of all isolates was identified as MDR, which higher than the rate of staphylococci MDR isolated from commensal specimens [21], and about 10 times more than the prevalence reported by Cavanagh et al. [2]. Second, *S. epidermidis* and other species (*S. warneri*, *S. cohnii*, *S. saprophyticus* and *S. hominis*) were all significantly associated with MDR, and all higher than *S. haemolyticus*. To our knowledge, this is the first comparison of MDR patterns among MR-CoNS collected from non-healthcare environments. However, studies with larger samples are needed to confirm these findings.

We found 11 *mecA* positive isolates that were not resistant to oxacillin and cefoxitin. This may be explained by: (a) not all *mecA* positive staphylococci are resistant to oxacillin due to the low expression of PBP2a causing the low levels of minimum inhibitory concentration (MIC) [40], and (b) MR-CoNS can be incorrectly characterized by cefoxitin disk diffusion (35). Additionally, 63.4% of all isolates in the present study carried the *qacA/B* gene. This prevalence was higher than the rate of *qacA/B* gene carried by CoNS isolated from surgical sites [37], nurses and the general population in Hong Kong [44].

According to previous studies, the *SCCmec* types in MR-CoNS are more diverse than in MRSA [11]. Hanssen et al. [8] revealed that staphylococcal strains from the same geographical region carry identical *ccr* genes and differ from sequences of strains from other regions. This agreement supports the evidence of horizontal *SCCmec* gene transfer among staphylococcal strains [9,46]. All 41 MR-CoNS isolates in our study were characterized for their *SCCmec* types, and we found that most of them (23/41) were assigned as untypeable. This result correlated with *S. hominis* isolates from blood that carried the high rate (82%) of untypeable *SCCmec* types [18]. These untypeable *SCCmec* were suspected to carry novel *SCCmec* types as described previously [24,45]. However, the bias may have occurred due to the use of primer sets developed for *S. aureus*. In addition to this result, 4/8 of *SCCmec* IVa and 1/1 of *SCCmec* IVd belonged to methicillin-resistant *S. epidermidis* (MRSE). This supported the description of Wisplinghoff et al. [39] and Du et al. [5] that found *SCCmec* IV in most MRSE strains.

In conclusion, the university environments such as library,

canteen, restrooms and computer room are the essential reservoirs of MR-CoNS. The species distribution of these isolates was similar to the strains isolated from clinical and commensal specimens. We demonstrated the high prevalence of quaternary ammonium resistance gene of these MR-CoNS and most of the isolates were multidrug resistant bacteria. This finding provided useful information to support disease prevention strategies against staphylococcal infections. People should be careful when touching these surfaces because they can be the carrier of these high-antibiotic resistance bacteria to other people. Hand washing activity should be usually practiced to eliminate this reservoir. 🇹🇭

Acknowledgement. This work was supported by a grant from the National Research Council of Thailand (R2560B064) to SS. RS was supported by the Royal Scholarship under Her Royal Highness Princess Maha Chakri Sirindhorn, 2015 of Naresuan University. We also acknowledge Dr. Keiichi Hiramatsu and Dr. Teruyo Ito for providing *SCCmec* type strains. Many thanks to Mr. Roy Morien of the Naresuan University Language Centre for his editing assistance and advice on English.

Competing interests. None declared.

References

1. Ardic N, Ozyurt M, Sareyyupoglu B, Haznedaroglu T (2005) Investigation of erythromycin and tetracycline resistance genes in methicillin-resistant staphylococci. *Int J Antimicrob Agent* 26:213-218
2. Cavanagh JP, Wolden R, Heise P, Esaiassen E, Klingenberg C, Aarag Fredheim EG (2016) Antimicrobial susceptibility and body site distribution of community isolates of coagulase-negative staphylococci. *APMIS* 124:973-978
3. Chaieb K, Zmantar T, Chehab O, Boucham O, Hasen AB, Mahdouani K, Bakhrouf A (2007) Antibiotic resistance genes detected by multiplex PCR assays in *Staphylococcus epidermidis* strains isolated from dialysis fluid and needles in a dialysis service. *Jpn J Infect Dis* 60:183
4. Chiang YC, Lu HC, Li SC, Chang YH, Chen HY, Lin CW, Tsen HY (2012) Development of PCR primers and a DNA microarray for the simultaneous detection of major *Staphylococcus* species using *groESL* gene. *Foodborne Pathog Dis* 9:249-257
5. Du X, Zhu Y, Song Y, Li T, Luo T, Sun G, Li M (2013) Molecular analysis of *Staphylococcus epidermidis* strains isolated from community and hospital environments in China. *PLoS One* 8:e62742
6. Eed, EM, Ghonaim MM, Hussein YM, Saber T, Khalifa AS (2015) Phenotypic and molecular characterization of HA-MRSA in Taif hospitals, Saudi Arabia. *J Infect Dev Ctries* 9:298-303
7. Friedman L, Wallar LE, Papadopoulos A (2015) Environmental risk fac-

- tors for community-acquired MRSA. National Collaboration Center for Environment Health. Retrieved from http://www.nccch.ca/sites/default/files/Environmental_Risk_Factors_MRSA_April_2015.pdf
8. Hanssen AM, Sollid JUE (2006) SCC*mec* in staphylococci: genes on the move. *FEMS Immunol Med Microbiol* 46:8-20
 9. Hanssen AM, Kjeldsen G, Sollid JUE (2004) Local variants of staphylococcal cassette chromosome *mec* in sporadic methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative staphylococci: evidence of horizontal gene transfer? *Antimicrob Agents Chemother* 48:285-296
 10. Hirota S, Sasaki T, Kuwahara-Arai K, Hiramatsu K (2011) Rapid and accurate identification of human-associated staphylococci by use of multiplex PCR. *J Clin Microbiol* 49:3627-3631
 11. Hisata K, Kuwahara-Arai K, Yamanoto M, Ito T, Nakatomi Y, Cui L, Kinoshita S (2005) Dissemination of methicillin-resistant staphylococci among healthy Japanese children. *J Clin Microbiol* 43:3364-3372
 12. Kittit T, Boonyonying K, Sitthisak S (2011) Prevalence of methicillin resistant *Staphylococcus aureus* among university students in Thailand. *Southeast Asian J Trop Med Public Health* 42:1498-1504
 13. Kohner P, Uhl J, Kolbert C, Persing D, Cockerill FR (1999) Comparison of susceptibility testing methods with *mecA* gene analysis for determining oxacillin (methicillin) resistance in clinical isolates of *Staphylococcus aureus* and coagulase-negative *Staphylococcus* spp. *J Clin Microbiol* 37:2952-2961
 14. Koksall F, Yasar H, Samasti M (2009) Antibiotic resistance patterns of coagulase-negative *Staphylococcus* strains isolated from blood cultures of septicemic patients in Turkey. *Microbiol Res* 164:404-410
 15. Lally RT, Ederer MN, Woolfrey BF (1985) Evaluation of mannitol salt agar with oxacillin as a screening medium for methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 22:501-504
 16. Loonen AJ, Jansz AR, Bergland JN, Valkenburg M, Wolffs PF, van den Brule AJ (2012) Comparative study using phenotypic, genotypic, and proteomics methods for identification of coagulase-negative staphylococci. *J Clin Microbiol* 50:1437-1439
 17. Lwanga SK, Lemeshow S (1991) Sample size determination in health studies: a practical manual. World Health Organization, Geneva, Switzerland
 18. Mendoza-Olazarán S, Morfin-Otero R, Rodríguez-Noriega E, Llacadiaz J, Flores-Treviño S, González-González GM, Garza-González E (2013) Microbiological and molecular characterization of *Staphylococcus hominis* isolates from blood. *PLoS One* 8:e61161
 19. Mishra SK, Rijal BP, Pokhrel BM (2013) Emerging threat of multidrug resistant bugs—*Acinetobacter calcoaceticus baumannii* complex and methicillin resistant *Staphylococcus aureus*. *BMC Res Notes* 6:1
 20. Mkrtychyan HV, Russell CA, Wang N, Cutler R (2013) Could public restrooms be an environment for bacterial resistomes? *PLoS One* 8:e54223
 21. Morgenstern M, Erichsen C, Hackl S, Mily J, Militz M, Friederichs J, Post V (2016) Antibiotic resistance of commensal *Staphylococcus aureus* and coagulase-negative staphylococci in an international cohort of surgeons: a prospective point-prevalence study. *PLoS One* 11:e0148437
 22. Osman, KM, Badr J, Al-Maary K, Moussa I, Hessain A, Girh Z, Saad A (2016) Prevalence of the antibiotic resistance genes in coagulase-positive-and negative-*Staphylococcus* in chicken meat retailed to consumers. *Front Microbiol* 7:1846
 23. Otto M (2013) Coagulase-negative staphylococci as reservoirs of genes facilitating MRSA infection. *Bioessays* 35:4-11
 24. Panda S, Jena S, Sharma S, Dhawan B, Nath G, Singh DV (2016) Identification of novel sequence types among *Staphylococcus haemolyticus* isolated from variety of infections in India. *PLoS One* 11:e0166193
 25. Rahman A, Hosain M, Mahmud C, Paul S, Sultana S, Haque N, Kubayashi N (2012) Species distribution of coagulase negative staphylococci isolated from different clinical specimens. *Mymensingh Med J* 21:195-199
 26. Ryffel C, Tesch W, Birch-Machin I, Reynolds PE, Barberis-Maino L, Kayser FH, Berger-Bächi B (1990) Sequence comparison of *mecA* genes isolated from methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Gene* 94:137-138
 27. Sasaki T, Tsubakishita S, Tanaka Y, Sakusabe A, Ohtsuka M, Hirota S, Hiramatsu K (2010) Multiplex-PCR method for species identification of coagulase-positive staphylococci. *J Clin Microbiol* 48:765-769
 28. Schmidt T, Kock MM, Ehlers MM (2015) Diversity and antimicrobial susceptibility profiling of staphylococci isolated from bovine mastitis cases and close human contacts. *J Dairy Sci* 98:6256-6269
 29. Seshadri D, Sethuraman G (2016) Gram positive bacterial infections. In Singal, A, & Grover C. *Comprehensive approach to infections in dermatology*. Jaypee brothers Medical Pub, London, UK
 30. Shah MM, Iihara H, Noda M, Song S.X, Nhung P.H, Ohkusu K, Ezaki T (2007) *dnaJ* gene sequence-based assay for species identification and phylogenetic grouping in the genus *Staphylococcus*. *Int J Syst Evol Microbiol* 57:25-30
 31. Shobha K, Rao P, Thomas J (2005) Survey of *Staphylococcus* isolates among hospital personnel, environment and their antibiogram with special emphasis on methicillin resistance. *Indian J Med Microbiol* 23:186
 32. Shome B, Das Mitra S, Bhuvana M, Krithiga N, Velu D, Shome R, Rahman H (2011) Multiplex PCR assay for species identification of bovine mastitis pathogens. *J Appl Microbiol* 111:1349-1356
 33. Soge OO, Meschke JS, No DB, Roberts MC (2009) Characterization of methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative *Staphylococcus* spp. isolated from US West Coast public marine beaches. *J Antimicrob Chemother* 64:1148-1155
 34. Strommenger B, Kettlitz C, Werner G, Witte W (2003) Multiplex PCR assay for simultaneous detection of nine clinically relevant antibiotic resistance genes in *Staphylococcus aureus*. *J Clin Microbiol* 41:4089-4094
 35. Swenson JM, Tenover FC, Group CDS (2005) Results of disk diffusion testing with cefoxitin correlate with presence of *mecA* in *Staphylococcus* spp. *J Clin Microbiol* 43:3818-3823
 36. Tangchaisuriya U, Yotpanya W, Kittit T, Sitthisak S (2014) Distribution among Thai children of methicillin resistant *Staphylococcus aureus*. *Southeast Asian Trop Med Public Health* 42:149-156
 37. Temiz M, Duran N, Dura G G, Eryilmaz N, Jenedi K (2014) Relationship between the resistance genes to quaternary ammonium compounds and antibiotic resistance in staphylococci isolated from surgical site infections. *Sci Monit* 20:544
 38. Widerström M, Wiström J, Edebro H, Marklund E, Backman M, Lindqvist P, Monsen T (2016) Colonization of patients, healthcare workers, and the environment with healthcare-associated *Staphylococcus epidermidis* genotypes in an intensive care unit: a prospective observational cohort study. *BMC Infect Dis* 16:743

39. Wisplinghoff H, Rosato AE, Enright MC, Noto M, Craig W, Archer GL (2003) Related clones containing SCCmec type IV predominate among clinically significant *Staphylococcus epidermidis* isolates. *Antimicrob Agents Chemother* 47:3574-3579
40. Xu Z, Mkrtychyan HV, Cutler RR (2015) Antibiotic resistance and *mecA* characterization of coagulase-negative staphylococci isolated from three hotels in London, UK. *Front Microbiol* 6:947
41. Youn JH, Park YH, Hang'ombe B, Sugimoto C (2014) Prevalence and characterization of *Staphylococcus aureus* and *Staphylococcus pseudintermedius* isolated from companion animals and environment in the veterinary teaching hospital in Zambia, Africa. *Comp Immunol Microbiol Infect Dis* 37:123-130
42. Zell C, Resch M, Rosenstein R, Albrecht T, Hertel C, & Götz F (2008) Characterization of toxin production of coagulase-negative staphylococci isolated from food and starter cultures. *Int J Food Microbiol* 127:246-251
43. Zhang K, McClure JA, Elsayed S, Louie T, Conly JM (2005) Novel multiplex PCR assay for characterization and concomitant subtyping of staphylococcal cassette chromosome *mec* types I to V in methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 43:5026-5033
44. Zhang M, O'Donoghue M, Ito T, Hiramatsu K, Boost M (2011) Prevalence of antiseptic-resistance genes in *Staphylococcus aureus* and coagulase-negative staphylococci colonising nurses and the general population in Hong Kong. *J Hosp Infect* 78:113-117
45. Zong Z, Lü X (2010) Characterization of a new SCCmec element in *Staphylococcus cohnii*. *PLoS One* 5:e14016
46. Zong Z, Peng C, Lü X (2011) Diversity of SCCmec elements in methicillin-resistant coagulase-negative staphylococci clinical isolates. *PLoS One* 6:e20191

