

Stanley Maloy*
Moselio Schaechter

Center for Microbial Sciences,
San Diego State University,
San Diego, California, USA

Received 10 January 2006
Accepted 6 February 2006

*Corresponding author:
S. Maloy
Center for Microbial Sciences
San Diego State University
5500 Campanile Drive
San Diego, CA 92182-4614, USA
Tel. 1- 619-5947123. Fax 1- 619-5945676
Email: smaloy@sciences.sdsu.edu

The era of microbiology: a Golden Phoenix

Summary. The discoveries over the last decade have demonstrated that microbiology is a central scientific discipline with practical applications in agriculture, medicine, bioremediation, biotechnology, engineering, and other fields. It is clear that the roles of microbes in nature are so diverse that the process of mining this genetic variation for new applications will continue long into the future. Moreover, the rapid rate of microbial evolution ensures that there will be no permanent solution to agricultural, medical, or environmental problems caused by microbes. These problems will demand a continual stream of creative new approaches that evolve along with the microbes. Thus, the excitement of this field will continue long into the future. However, these opportunities and imperatives demand a deep understanding of basic microbial physiology, genetics, and ecology. Major challenges that lay ahead are to impart the broad training needed to entice and enable the next generation of microbiologists, and to educate the public and government representatives about the continued and critical importance of this field for health and the economy. [*Int Microbiol* 2006; 9(1):1-7]

Key words: development of microbiology · microbial ecology · microbial cell biology · integrative microbiology

Introduction

The development of powerful new technologies in the post-World War II period led to a revolution in biology. The introduction of genetic tools such as transposons, coupled with the ability to clone genes and determine DNA sequences, and the subsequent explosion of techniques based upon these methods made it possible to dissect the molecular genetics of essentially any organism. These developments allowed us to answer questions that were previously deemed unfathomable, and had a major impact on every discipline of biology. The study of microbes has particularly profited from these developments. It was previously only feasible to study details of microbial physiology of pure cultures growing in the laboratory. However, it is now possible to directly characterize organisms growing in

the natural environment, and to monitor the physiology of single cells under defined conditions. Advances in microscopy, nanotechnology, and structural biology allow the analysis of the underlying chemical mechanisms at a deeper and deeper level. In addition to enhancing our understanding of the biological world, these new developments have many practical applications. The ability to study microbes in their own environment rather than the test tube, coupled with the ability to dissect the molecular details of microbial cells has made microbial biology pivotal to nearly all disciplines of science. To illustrate this point, we will discuss the key stages in development of microbiology and how new developments have led to what we refer to as the third Golden Age of microbiology [37].

The first Golden Age

The existence of microbes and their role in disease was guessed by ancient Greeks and demonstrated convincingly by a number of studies in the 17th, 18th, and early 19th centuries.

This article is based on the invited lecture given by S. Maloy, president of the American Society for Microbiology, at the 20th National Congress of the Spanish Society for Microbiology on September 19, 2006, in Cáceres, Spain.

Despite what seems today like compelling evidence, most of these studies did not gain universal acceptance and were the subject of controversy. In the second half of the 19th century, microbiology made a grandiose entry, sweeping away doubts about the existence and importance of microbes. Within a 20-year period, the main bacterial etiological agents of disease in humans and animals were discovered, and the field of immunology was developed, leading to vaccines and serological tests. The discovery of the microbial etiology of major infectious diseases led in turn to accurate diagnoses and attempts at prevention and cure. Several vaccines used today stem from those developed by early microbiologists. The importance of microbes in the cycles of nature was elucidated and strain selection was applied for industrial purposes. The intellectual developments and practical applications of this first Golden Age propelled microbiology into becoming one of the main branches of biology and medicine (Table 1).

The second Golden Age

In the first half of the 20th century, microbiological research continued at an ever-increasing pace and many of the missing details in biochemical, medical, and environmental microbiology were uncovered. However, microbiology became increasingly fragmented. The unity that Pasteur originally brought to the discipline by applying similar approaches to medical and environmental microbiology was replaced by specialized subdisciplines. During this period, medical microbiologists and immunologists were mainly working at the organism level of host and parasite. Environmental microbiologists, on the other hand, were focused on chemical processes. Thanks to the pioneering work of the Russian Sergei Winogradsky and

Table 1. The first Golden Age of microbiology

Concepts	
Bacterial physiology	
Methods for cultivation and observation of bacteria	
Bacterial nutrition	
Bacterial classification based upon phenotypes	
Role of bacteria in cycles of nature - chemosynthesis and photosynthesis	
Medical microbiology	
Microbes as agents of acute disease in animals and plants	
Viruses	
Immunology	
Phagocytosis	
Antibodies	
Applications	
Clinical identification of microbes	
Antimicrobial chemotherapy	
Vaccines	
Industrial fermentations	

Table 2. The second Golden Age of microbiology

Concepts	
Bacterial genetics	
DNA as genetic material and its structure	
Genetic code	
Mechanism of gene expression	
Regulation of gene expression	
Transposons	
Bacterial physiology	
Membrane transport and electrochemical gradients	
Cellular immunology	
Applications	
Genetic engineering	
Nucleic acid and protein sequencing	
Microbial classification based upon genotypes	
Monoclonal antibodies	

the Dutch Martinus Beijerinck and Albert Kluyver, environmental microbiology became a branch of comparative biochemistry, emphasizing the unity of biochemical processes in all living organisms. Unfortunately, the great gains of this period in microbial biochemistry and geochemistry did not find a ready parallel in medical microbiology. Many advances were made but only a few startling breakthroughs emerged in this field. Microbial genetics, for most purposes, was still to be discovered, meaning that many cellular phenomena remained undecipherable. Although individual aspects of the field made important strides during this period, microbiology was not seen as a dominant science. All this changed as abruptly as the first Golden Age with the advent of molecular genetics.

The discovery of biochemical genetics and of genetic exchange mechanisms in bacteria and viruses ushered in a new era of stellar advances. These discoveries led to modern concepts of the gene and the biochemical basis of genetics, the understanding of how genetic information flows from nucleic acids to proteins, the regulation of gene expression, and how complex structures such as bacteriophages are assembled. These breakthroughs led to a paradigm shift. At that time, anyone who wanted to do modern science, mindful of it or not, had to become a microbiologist. The incipient science of molecular biology was spawned by the use of microbes and, consequently microbial science was once again recognized as a fundamental scientific discipline (Table 2).

The third Golden Age

Soon after the discoveries in molecular genetics were made, investigators studying eukaryotic systems adopted the new tools for their own use. For those systems that can be manip-

ulated with nearly the same ease as bacteria, such as yeast, the transition was straightforward. Ultimately, other systems that lack the ease of manipulation and the speed of growth of bacteria benefited even more from DNA cloning and sequencing. As the methodological playing field became more even, eukaryotic science blossomed, eventually overshadowing the work with bacteria and phages. As funding for basic microbiology became increasingly difficult to obtain in the United States and elsewhere, many aspects of this science became neglected and were left to a few stalwarts. Although modern genetic techniques appropriate to “higher” organisms were gradual in coming, they promised the genetic analysis of such biological phenomena as cancer and differentiation. In a short time, enormous progress was made along these lines. By the last decades of the 20th century, the study of prokaryotes played distinct second fiddle. In addition, the early success of antibiotics and their promise dampened the impetus for the study of microbial pathogens. William Stewart, the Surgeon General of the United States, echoed the conviction of many scientists when he said in 1967 that it was “time to close the book on infectious diseases, declare the war against pestilence won, and shift national resources to such chronic problems as cancer and heart disease”. Not surprisingly, microbiology lost its luster in the eyes of many, leading to the wholesale closure of microbiology departments and diminishing funding opportunities for researchers in this field. Despite rising from the ashes of the first Golden Age, the success of microbiology was responsible for its own demise.

All this has changed in the last 20 years, restoring the eminence of microbial science. Discoveries during this time have led to the understanding that microbes are essential for processes that impinge on nearly every aspect of our planet. It is now clear that microbial biology plays a unique and fundamental role in essentially every field of science. These insights have largely come from the study of microbes at the two ends of the spectrum—from the broad perspective of microbes in natural environments to the focused perspective of the molecular mechanisms operating in microbial cells (Table 3).

Microbial ecology and population biology

As eloquently argued by paleontologist Stephen Jay Gould [19], the 21st century will be the era of the microbe.

“We live now in the ‘Age of Bacteria.’ Our planet has always been in the “Age of Bacteria,” ever since the first fossils—bacteria, of course—were entombed in rocks more than 3 billion years ago. On any possible, reasonable or fair criterion, bacteria are—and always have been—the dominant forms of life on Earth. Our failure to grasp this most

Table 3. The third Golden Age of microbiology

Concepts

- Genomics and evolution
 - Extent of horizontal gene exchange
 - Diversity in microbial populations
 - Emerging infectious diseases
- Microbial ecology
 - Identification of uncultivated microbes
 - Role of microbes in modulating host development
 - Interactions between microbes
- Microbial physiology
 - Mechanisms of signal transduction
 - Global regulatory mechanisms
 - Interactions between proteins
 - Metabolic networks
- Innate immunity
- Role of microbes in chronic diseases

Applications

- Identification of uncultivated microbes
- New methods for the rapid identification of microbes
- New targets for antimicrobial therapies
- Rational development of probiotics
- Metabolic engineering
- Use of microbes as nanomachines
- Use of microbes for bioremediation

evident of biological facts arises in part from the blindness of our arrogance but also, in large measure, as an effect of scale. We are so accustomed to viewing phenomena of our scale—sizes measured in feet and ages in decades—as typical of nature.”

Despite their small size, microbes play an enormous role in the cycle of matter and in the metabolism of this planet. With the discovery of the huge microbial biota in subsoil fissures and better estimates of microbial life in the oceans, the microbial biota on Earth is thought to exceed in weight all other living things combined. Bacteria alone account for 50% of the biomass of carbon and over 90% of the biomass of nitrogen and phosphorus on our planet. At least as much photosynthesis is carried out by marine microbes as by terrestrial plants [40]. It has been estimated that if the action of microbes on the nitrogen cycle were to cease, the amount of nitrogen available to plants would become too low to sustain life within about one week (David Lipson, personal communication). Microbes are also critical for assimilating dissolved organic material into the particulate organic matter used by eukaryotes [3]. Microbes are responsible for enormous geochemical activities, such as the formation of carbon and silica deposits and the shaping of limestone caverns [13]. Microorganisms can be found in unexpected niches, such as thermal ocean vents and fissures in deep rocks [44]. Such lithotrophic organisms support the existence of biomes that are not ultimately dependent on the sun or atmospheric oxygen, expanding the scope of microbial ecolo-

gy and our understanding of the role of microbes in the cycles of matter in nature, but also raising new questions about microbial physiology [48].

These insights come largely from the ability to study microbes in nature. Once a specialized branch of microbiology, microbial ecology now occupies center stage. Thanks to cloning techniques and PCR, it became possible to break out of the confines of model organisms grown in the laboratory to study the physiology and genetics of microbes in the natural environment. This includes species that cannot yet be cultured in the laboratory, estimated to comprise over 99% of the total microbial population [43]. It is now possible to approach questions such as who lives where, who does what, and who is phylogenetically related to whom. This has greatly expanded our awareness of the diversity of types and total numbers of microbes, even embracing a new Domain of life, the *Archaea*, a division that we did not know existed during the two previous golden ages.

As the extent of microbial diversity has become appreciated, many microbiologists have questioned the concept of non-cultivability—a concept that simply reflects experimental failure rather than a verifiable conclusion. This rationale has driven the development of new methods to culture species previously believed to be uncultivable [42]. We can expect many more of these “uncultivable” microbes to be grown in the laboratory, facilitating studies on their physiology and potentially allowing them to be harnessed for practical applications.

These approaches have led to an increased appreciation that microbes in nature tend to live in communities, some with their own kin, others in consortia with different microbes. This awareness dictates that the study of microbes must be extended from the laboratory to the natural communities. In fact, pathogenic microbes are increasingly being investigated in their natural habitat. Clever genetic approaches have been developed to identify genes that are selectively expressed in a particular environment. These approaches have many applications, for example identifying gene products that are essential for a pathogen to survive in a host or in the outside environment [10]. This ecological approach has led to great strides in the identification of novel molecular mechanisms of cell–cell interactions. An example is the Type III secretion system, a device used by bacteria to “inject” proteins directly into host cells and which is turned on by contact with host cells [30]. This implausible-sounding mechanism is shared by a surprisingly large number of pathogens of vertebrates, insects, worms, and plants [6]. Like earlier studies on microbial physiology that led to the concept of the unity of biochemistry, these findings suggest that microbe–host interactions may share common mechanisms despite the type of host or whether the interaction is mutualistic or parasitic.

Recent studies on microbe–host interactions indicate that microbes influence the host in ways we never imagined previously. For example, the presence of the normal microbial biota in the intestine of mice promotes proper development of the vascular tissue as well as host immunity to invading pathogens [4,41]. These discoveries provide novel insights into beneficial microbe–host interactions that may lead to rational design of probiotics. However, we are also beginning to learn that microbes can cause chronic diseases that were previously thought to be due to genetics or environment—ailments such as ulcers and stomach cancer caused by *Helicobacter pylori*, arthritis caused by *Borrelia burgdorferi*, and many other diseases [7]. Epidemiological evidence coupled with sensitive assays for bacteria based upon the environmental PCR approaches hint that the role of microbes in causing chronic diseases is pervasive. These studies have also led to a deeper understanding of innate immunity.

Genomics and evolution

In part because of their small size, the genomes of many prokaryotes have been sequenced. The number of completed sequences includes not only representatives of major groups but also multiple strains of many species. Not surprisingly, this richness of data has spawned a whole industry intended to “mine” the genomic information for practical uses [14,25]. Genome sequences have provided important insights into evolution. One surprise has been the demonstration of the extent and impact of lateral gene transfer. Genomic islands, sets of genes a dozen or more in number, appear in unrelated microbes and endow them with new functions such as a virulence trait or a novel metabolic capacity [12]. Often these functions are encoded on mobile genetic elements that can readily move between bacteria [15]. For example, many bacterial toxins are encoded on bacteriophages that can integrate into the bacterial genome [46]. Thus, microbes break out of the confines of sexual recombination by the promiscuous transfer of genes between organisms. Most of the DNA so transferred is lost, but some is retained to alter the genotype and phenotype of recipients [22]. The extent and impact of lateral gene transfer has made us reevaluate the concept that evolution can be linearly traced along separated branches of a tree. Instead, the branches of the tree are laterally connected to one another. Thus, the ancestry of an organism may not be simply monophyletic, but can result from multiple exchanges between distinct genomes. Such interactions are not limited to one Domain of life, but can also jump between *Bacteria* and *Archaea* or *Eukarya* [2]. This lateral gene exchange seriously compromises the concept of species in microbiology [17].

Lateral gene transfer does not tell the whole story of microbial evolution. Genome sequences have also revealed the importance of point mutations (SNPs) in bacterial evolution [47]. The accumulation of pseudogenes seems to play an important role in restricting a microbe to a particular host or environmental niche [29]. These discoveries have an important impact on our understanding of how new emerging diseases arise, but we still do not know how often these events occur in nature.

Microbial cell biology

An old dictum argued that bacteria are just bags of enzymes. It is becoming increasingly clear that within the small confines of one cubic millimeter or so, there is a sophisticated and unexpected compartmentalization [5]. Cellular components act in dynamic fashion and a surprising molecular choreography has been unveiled via fluorescence microscopy and other techniques that allow us to visualize molecules in action [23,35]. Some macromolecules localize to specific sites, others move around the cell interior along established paths. In nearly all prokaryotes, cell division depends on the formation at mid-cell of a constricting ring that consists of the tubulin-like protein FtsZ [26]. Other proteins, called Min, bounce from one pole of a rod-shaped bacterium to the other, helping to determine the division site. Rod-shaped bacteria also have the equivalent of actin and some that of intermediate filaments [16,34]. Thus, at least some of the eukaryotic cytoskeletal proteins were prokaryotic inventions that long preceded emergence in eukaryotes.

Microbial symbionts and pathogens make extensive uses of host cell machinery, such as cell surface receptors, cell signaling pathways, and the cytoskeleton [1]. An example is the appropriation of host cell actin by *Escherichia coli* to promote binding to intestinal cells [39] or by *Shigella* and *Listeria* to facilitate direct passage from one host cell to another [18], thereby avoiding exposure to the host immune system. In addition to providing insights into pathogenesis, these processes provide tools to understand the structure and function of eukaryotic cells. Nowadays, pathogenic microbiologists are being invited with increasing frequency to deliver papers at cell biology meetings. As in the old days, when the proper study of bacteria was via bacteriophages, it seems increasingly evident that studying bacteria–host interactions provides unique tools for studying many aspects of eukaryotic cell biology. These conceptual advances have many practical applications. The increasing problem of antibiotic resistance, the burden of emerging infectious diseases, and the threat of bioterrorism require improved ways of rapidly

detecting pathogenic agents, preventing their transmission, and effectively treating infected humans, other animals, and plants. These problems demand a continuous search for new antimicrobial drugs to stay one step ahead of the evolution of microbial resistance [11], as well as the development of effective new vaccines to treat emerging diseases [21]. Both of these applications are increasingly relying on a detailed understanding of genomics, structural biology, molecular biology, and microbial physiology. For example, understanding the details of genetic recombination has led to specific peptide inhibitors of this essential cellular function [20]. Likewise, understanding of protein structure and function provides new approaches for designing effective vaccines.

Microbiology is having an increasing impact in other areas as well. Microbes are playing an increasing role in industrial production of pharmaceuticals and other compounds [27]. There has already been considerable success in mining the genomes of cultivated and uncultivated microorganisms for enzymes with desirable physical and chemical properties [45]. This approach has already brought to market several hydrolytic enzymes with enhanced thermal stability and targeted substrate specificities. But there are many other opportunities for industrial microbiology. For example, fermentation processes have the potential to decrease our reliance on fossil fuels [24], and bacterial systems are being developed as nano-biosensors [9], nano-batteries [38], and nano-computers [49]. An infusion of funds in these areas will almost certainly lead to significant advances, but note that these practical applications ultimately depend on a strong foundation of basic scientific research and a deep understanding of the details of bacterial physiology.

Integrative microbiology

Microbiology has become more unified. The fields of microbial physiology, microbial genetics, microbial ecology, and microbial pathogenesis are no longer sovereign domains with their own tools and dialects. Nowadays, a marine microbiologist can easily talk to a microbiologist studying human pathogens, and a food microbiologist can converse effortlessly with a scientist studying microbial evolution. These interactions are not simply because scientists working on different problems share genomic databases and molecular tools—the interactions are driven by the realization that microorganisms use similar mechanisms to accomplish diverse functions. Discoveries in microbial pathogenesis have a direct impact on the field of microbial ecology, and an understanding of microbial ecology is critical for understanding transmission of pathogens. Common molecular mechanisms of adhesion to surfaces, quorum sensing, signal transduction,




Fig. 1. For many years microbiology has had a major impact on the disciplines shown on the left side of the figure. However, as we have learned more about the impact of microbes on the environment, the roles of microbes in causing diseases previously attributed to genetics or environment, and the applications of microbes in nanotechnology, microbiology has assumed an important role in essentially all disciplines of science, including the disciplines shown on the right side of the figure.

responding to environmental stresses, or injecting proteins directly into host cells are found in a wide variety of microbes that live in a broad spectrum of environmental niches, from humans to the ocean to sulfurous hot springs. As a result, the subdisciplines of microbiology are no longer isolated fields of study (Fig. 1).

In addition to its roles in agriculture and medicine, microbiology has also become an integral part of other scientific disciplines. The role of microbes in geochemistry has made microbiology a fundamental aspect of geology [31]. The use of microbes in nanotechnology has brought microbiology to engineering [36], and physics [28]. Microbial processes profoundly influence climate and weather, integrating microbiology into the field of atmospheric sciences [33]. Microbes are increasingly used to make complex molecules with chiral properties that cannot be synthesized in a test tube, making microbiology an invaluable assistant in chemistry [32]. And recent discoveries indicate that microbes also play important roles in determining animal behavior, bringing microbiology to the realm of psychology [8]. Hence, microbiology has become an integrative science that simultaneously adopts the tools of diverse scientific disciplines and impacts diverse scientific disciplines.

References

- Alonso A, García-del Portillo F (2004) Hijacking of eukaryotic functions by intracellular bacterial pathogens. *Int Microbiol* 7:181-191
- Andersson JO (2005) Lateral gene transfer in eukaryotes. *Cell Mol Life Sci* 62:1182-1197
- Azam F, Smith DC, Stewart GF, Hagström Å (1993) Bacteria-organic matter coupling and its significance for oceanic carbon cycling. *Microb Ecol* 28:167-179
- Backhed F, Ley RE, Sonnenburg JL, Peterson DA, Gordon JI (2005) Host-bacterial mutualism in the human intestine. *Science* 307:1915-1920
- Bravo A, Serrano-Heras G, Salas M (2005) Compartmentalization of prokaryotic DNA replication. *FEMS Microbiol Rev* 29:25-47
- Büttner D, Bonas U (2003) Common infection strategies of plant and animal pathogenic bacteria. *Curr Opin Plant Biol* 6:312-319
- Cabone K, Luftig R, Buckley M (2005) Microbial triggers of chronic human disease. ASM Press, Washington DC
- Callahan GN (2002) Madness. *Emerg Infect Dis* 8:998-1002
- Cheng Vollmer A, Van Dyk TK (2004) Stress responsive bacteria: biosensors as environmental monitors. *Adv Microb Physiol* 49:131-174
- Chiang SL, Mekalanos JJ, Holden DW (1999) In vivo genetic analysis of bacterial virulence. *Annu Rev Microbiol* 53:129-154
- Courvalin P (2005) Antimicrobial drug resistance: "Prediction is very difficult, especially about the future". *Emerg Infect Dis* 11:1503-1506
- Dobrindt U, Hochhut B, Hentschel U, Hacker J (2004) Genomic islands in pathogenic and environmental microorganisms. *Nat Rev Microbiol* 2:414-424
- Engel A, Stern L, Bennett P (2004) Microbial contributions to cave formation: New insights into sulfuric acid speleogenesis. *Geology* 32:369-372
- Eyers L, George I, Schuler L, Stenuit B, Agathos SN, El Fantroussi S (2004) Environmental genomics: exploring the unmined richness of microbes to degrade xenobiotics. *Appl Microbiol Biotechnol* 66:123-130
- Frost LS, Leplae R, Summers AO, Toussaint A (2005) Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol* 3:722-732
- Gerdes K, Moller-Jensen J, Ebersbach G, Kruse T, Nordström K (2004) Bacterial mitotic machineries. *Cell* 116:359-366
- Gevers D, Cohan FM, Lawrence JG, Spratt BG, Coenye T, Feil EJ, Stackebrandt E, Van de Peer Y, Vandamme P, Thompson FL, Swings J (2005) Opinion: Re-evaluating prokaryotic species. *Nat Rev Microbiol* 3:733-739
- Gouin E, Welch MD, Cossart P (2005) Actin-based motility of intracellular pathogens. *Curr Opin Microbiol* 8:35-45
- Gould SJ (1996) Planet of the bacteria, *Washington Post Horizon*, vol. 119, p 344

20. Gunderson CW, Segall AM (2006) DNA repair, a novel antibacterial target: Holliday junction-trapping peptides induce DNA damage and chromosome segregation defects. *Mol Microbiol* 59:1129-1148
21. Kaper J, Rappuoli R, Buckley M (2005) Vaccine development: current status and future needs. ASM Press, Washington DC
22. Lawrence JG, Hendrickson H (2005) Genome evolution in bacteria: order beneath chaos. *Curr Opin Microbiol* 8:572-578
23. Lim GE, Derman AI, Pogliano J (2005) Bacterial DNA segregation by dynamic SopA polymers. *Proc Natl Acad Sci USA* 102:17658-17663
24. Lin Y, Tanaka S (2006) Ethanol fermentation from biomass resources: current state and prospects. *Appl Microbiol Biotechnol* 69:627-642
25. Lorenz P, Eck J (2005) Metagenomics and industrial applications. *Nat Rev Microbiol* 3:510-516
26. Margolin W (2005) FtsZ and the division of prokaryotic cells and organelles. *Nat Rev Mol Cell Biol* 6:862-871
27. Mir J (2004) Industrial microbiology. A new challenge. *Int Microbiol* 7:81-82
28. Mitchell JG, Kogure K (2006) Bacterial motility: links to the environment and a driving force for microbial physics. *FEMS Microbiol Ecol* 55:3-16
29. Moran NA, Plague GR (2004) Genomic changes following host restriction in bacteria. *Curr Opin Genet Dev* 14:627-633
30. Mota LJ, Cornelis GR (2005) The bacterial injection kit: type III secretion systems. *Ann Med* 37:234-249
31. Newman DK, Banfield JF (2002) Geomicrobiology: how molecular-scale interactions underpin biogeochemical systems. *Science* 296:1071-1077
32. Patel RN (2003) Microbial/enzymatic synthesis of chiral pharmaceutical intermediates. *Curr Opin Drug Discov Devel* 6:902-920
33. Pedrós-Alió C, Simó R (2002) Studying marine microorganisms from space. *Int Microbiol* 5:195-200
34. Pogliano J (2004) DNA segregation by bacterial actin homologs. *Dev Cell* 6:3-4
35. Rubio A, Jiang X, Pogliano K (2005) Localization of translocation complex components in *Bacillus subtilis*: enrichment of the signal recognition particle receptor at early sporulation septa. *J Bacteriol* 187:5000-5002
36. Sara M, Pum D, Schuster B, Sleytr UB (2005) S-layers as patterning elements for application in nanobiotechnology. *J Nanosci Nanotechnol* 5:1939-1953
37. Schaechter M (2003) Integrative microbiology—the third Golden Age. *J Biosci* 28:149-154
38. Scholz F, Schroder U (2003) Bacterial batteries. *Nat Biotechnol* 21:1151-1152
39. Selbach M, Backert S (2005) Cortactin: an Achilles' heel of the actin cytoskeleton targeted by pathogens. *Trends Microbiol* 13:181-189
40. Shively JM, English RS, Baker SH, Cannon GC (2001) Carbon cycling: the prokaryotic contribution. *Curr Opin Microbiol* 4:301-306
41. Stappenbeck TS, Hooper LV, Gordon JI (2002) Developmental regulation of intestinal angiogenesis by indigenous microbes via Paneth cells. *Proc Natl Acad Sci USA* 99:15451-15455
42. Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA (2004) New strategies for cultivation and detection of previously uncultured microbes. *Appl Environ Microbiol* 70:4748-4755
43. Streit WR, Schmitz RA (2004) Metagenomics—the key to the uncultured microbes. *Curr Opin Microbiol* 7:492-498
44. Teske AP (2005) The deep subsurface biosphere is alive and well. *Trends Microbiol* 13:402-404
45. van den Burg B (2003) Extremophiles as a source for novel enzymes. *Curr Opin Microbiol* 6:213-218
46. Waldor M, Freidman D, Adhya S (2005) Phages: their role in bacterial pathogenesis and biotechnology. ASM Press, Washington DC
47. Weissman SJ, Moseley SL, Dykhuizen DE, Sokurenko EV (2003) Enterobacterial adhesins and the case for studying SNPs in bacteria. *Trends Microbiol* 11:115-117
48. Wood AP, Aurikko JP, Kelly DP (2004) A challenge for 21st century molecular biology and biochemistry: what are the causes of obligate autotrophy and methanotrophy? *FEMS Microbiol Rev* 28:335-352
49. Yokobayashi Y, Weiss R, Arnold FH (2002) Directed evolution of a genetic circuit. *Proc Natl Acad Sci USA* 99:16587-16591

La era de la microbiología: un Fénix de oro

Resumen. Los descubrimientos de la última década han demostrado que la microbiología no es sólo una disciplina científica con aplicaciones prácticas en agricultura, medicina, biorremediación, biotecnología, ingeniería y otros campos. Está claro que el papel de los microbios en la naturaleza es tan diverso que la explotación de esta variación genética para nuevas aplicaciones va a continuar en el futuro. Además, la rápida evolución microbiana asegura que no habrá soluciones permanentes para los problemas de la agricultura, la medicina o el medio causados por los microbios. Estos problemas tendrán que ser abordados con nuevos enfoques que evolucionen a la vez que los microorganismos. Con ello, la fascinación de esta disciplina se mantendrá en el futuro. Pero estas oportunidades e imperativos exigen un profundo conocimiento de la fisiología, la genética y la ecología microbianas básicas. Los principales retos son proporcionar una formación amplia a fin de atraer y capacitar a la siguiente generación de microbiólogos, y educar a la población y a los gobernantes para entender la importancia crucial de esta disciplina en la salud y en la economía. [*Int Microbiol* 2006; 9(1):1-7]

Palabras clave: desarrollo de la microbiología · ecología microbiana · biología celular microbiana · microbiología integral

A era da microbiologia: uma Fênix de ouro

Resumo. Os descobrimentos da última década demonstraram que a microbiologia não é só uma disciplina científica com aplicações práticas em agricultura, medicina, biorremedio, biotecnologia, engenharia e outros campos. Está claro que as funções dos micróbios na natureza são tão diversas, que a exploração desta variação genética para novas aplicações vai continuar no futuro. No entanto, a rápida evolução microbiana determina que não haverá soluções permanentes para os problemas da agricultura, a medicina ou o meio causados pelos micróbios. Estes problemas terão que ser abordados com novos enfoques que evoluam ao mesmo tempo que os micróbios. Com isso, a fascinação desta disciplina se manterá no futuro. Mas estas oportunidades e imperativos exigem um profundo entendimento da fisiologia, a genética e a ecologia microbianas básicas. Os principais desafios são proporcionar uma formação ampla com o fim de atrair e capacitar à seguinte geração de microbiologistas, e educar ao público e aos governantes para entender a importância crucial desta disciplina na saúde e na economia. [*Int Microbiol* 2006; 9(1):1-7]

Palavras chave: desenvolvimento da microbiologia · ecologia microbiana · biologia celular microbiana · microbiologia integradora