

Contribution of the FeoB transporter to *Streptococcus suis* virulence

Jesús Aranda,¹ Pilar Cortés,¹ Maria E. Garrido,¹ Nahuel Fittipaldi,²
Montserrat Llagostera,¹ Marcelo Gottschalk,² Jordi Barbé^{1*}

¹Department of Genetics and Microbiology, Autonomous University of Barcelona, Bellaterra, Barcelona, Spain. ²Research Centre for Swine Infectiology, Faculty of Veterinary Medicine, University of Montréal, St-Hyacinthe, Québec, Canada

Received 11 March 2009 · Accepted 15 May 2009

Summary. The contribution of iron transporter systems encoded by *feo* genes to the pathogenic traits of streptococci is largely unknown, despite the fact that those systems are required for the full virulence of several gram-negative bacterial species. In this work, we show that the swine pathogen and zoonotic agent *Streptococcus suis* has a *feoAB* operon similar to that encoding an iron transporter system in *Escherichia coli*. Electrophoretic mobility assays and transcriptional analyses confirmed that the expression of *S. suis* *feo* genes is under the negative control of the ferric uptake regulator (Fur) protein. In vivo trials in mice using a *feoB* defective mutant strain were carried out to investigate the contribution of this gene to the virulence of *S. suis*. The results showed that the median lethal dose (LD₅₀) of the mutant was approximately 10-fold higher than that of the wild-type parent strain. These data suggest that the Feo metal transporter plays a significant role in streptococcal infectious disease. This is in contrast to previous results reported for this same gene in other gram-positive bacterial species. [Int Microbiol 2009; 12(2):137-141]

Keywords: *Streptococcus suis* · operon *feo* · iron uptake · bacterial virulence · streptococcal diseases

Introduction

Streptococcus suis is a major pathogen of swine that is responsible for severe economic losses in the pork industry [13]. In addition, this bacterium is an emerging zoonotic agent, considered to be the primary cause of adult meningitis in Vietnam and the second highest cause of this disease in Thailand [17,26]. Moreover, in a recent outbreak in China, more than 200 human cases of *S. suis* infection, resulting in 39 deaths, were reported [12]. In both pigs and humans, the main clinical manifestations of *S. suis* infection are septicemia and meningitis [12].

Iron is an essential cation for living organisms, although its availability in the body fluids of host organisms is

extremely low [1]. To cope with this scarcity, most bacterial pathogens have evolved many different iron uptake systems [30]. In *Escherichia coli*, for example, the Fe²⁺ transport system is encoded by *feo* genes, which act in concert to maintain cellular iron supplies under anoxic conditions [15,25].

In *Escherichia coli*, the *feoABC* operon encodes three predicted proteins: FeoA, a small 75-residue hydrophilic protein of unknown function; FeoB, a large 773-residue protein with an integral membrane domain likely to act as the ferrous permease; and FeoC, a small 78-residue hydrophilic protein that has been suggested to function as a transcriptional repressor, so far only associated with gammaproteobacterial *feo* systems [7,20]. FeoB homologs are found in both gram-negative and gram-positive bacteria [20]. Interestingly, it has been shown that these homologs contribute to the virulence of gram-negative organisms, such as *E. coli* [25], *Helicobacter pylori* [29], *Campylobacter jejuni* [20], *Porphyromonas gingivalis* [8], and *Legionella pneumophila* [22]. However, FeoB involvement in the virulence of gram-positive microorganisms has been analyzed only in *Listeria monocytogenes*.

*Corresponding author: J. Barbé
Departament de Genètica i Microbiologia
Facultat de Biociències, Universitat Autònoma de Barcelona
08193 Bellaterra, Spain
Tel. +34-935811837. Fax + 34-935812387
E-mail: jordi.barbe@uab.cat

In contrast to previous reports in gram-negative bacteria, the virulence of *L. monocytogenes feoB* mutants in mice is not affected [14].

To our knowledge, the contribution of *feo* genes to the virulence of members of the *Streptococcus* genus has never been reported. In this work, *in silico* analysis was used to identify a *feoAB* operon in *S. suis*. We show that this operon is under the control of the ferric uptake regulator Fur, and that the product of the *feoB* gene is a significant virulence factor for *S. suis*.

Materials and methods

Bacterial strains, plasmids, and molecular techniques. The bacterial strains and plasmids used in this work are listed in Table 1. *Escherichia coli* strains were grown in Luria-Bertani (LB) broth or agar [19]. When necessary, ampicillin (50 µg/ml), chloramphenicol (34 µg/ml), and 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG; Roche) were added to the growth media. *Streptococcus suis* was grown in Todd-Hewitt medium (TH; Difco). Growth under metal-depleted conditions was achieved by the addition of the metal chelator 2,2'-dipyridyl (DPD; Sigma), at a final concentration of 100 µM, to TH broth 30 min before the inoculation of *S. suis* cells. When required, spectinomycin (100 µg/ml) and chloramphenicol (5 µg/ml) were added to TH medium. DNA extraction, plasmid mini-preparations, cloning, and transformation were carried out using standard procedures [24].

Construction of the *Streptococcus suis feoB* mutant. *Streptococcus suis* UA5003 mutant (*feoB::spc*) was obtained by insertional mutagenesis using allelic replacement. Briefly, the *feoB* gene of *S. suis* strain P1/7 was PCR-amplified using Pwo DNA polymerase (Roche) and primers 1638upSma and 1638dwSma (Table 2), which introduce *Sma* I restriction sites at both ends. The amplicon was cloned into the pGEM-T vector (Promega). Afterwards, the entire construction was amplified by overlap extension PCR with primers 1638intUP and 1638intDW (Table 2), which

anneal to an internal region of the *feoB* gene. A spectinomycin (*spc*) cassette released from pR412 [18] was treated to generate blunt ends and then ligated into the purified PCR product using T4 DNA ligase (Roche). After the resulting *feoB::spc* fragment had been digested with *Sma* I, the inactivated gene was cloned into the thermosensitive suicide vector pSET5s [28], giving rise to plasmid pSET5s*feoB::spc*. This vector was propagated in *E. coli* DH5α cells (Clontech) according to the manufacturer's instructions, purified, and introduced into *S. suis* strain P1/7 by electroporation [27]. Allelic replacement procedures were carried out as described previously [3]. Gene replacement in candidate clones was confirmed by PCR and by sequencing of the region in the resulting mutant using primers SecAup2 and SecAup3 (Macro-Gen Sequencing Service) (Table 2).

Virulence assays. All animal experiments were approved by the UAB Animal Ethics Committee. Female BALB/cAnNHsd mice (8-week old) obtained from Harlan Iberica (Barcelona, Spain) were used for virulence assays [2]. Four groups of three mice were used for each of the wild-type (WT) P1/7 or mutant UA5003 (*feoB::spc*) strains. Animals were injected intraperitoneally as described [11] with 0.1 ml of serial 10-fold dilutions of bacteria in suspensions of TH broth, supplemented with 10% inactivated bovine serum (Invitrogen). The concentrations of the original bacterial suspensions were determined by the plate count method. Animals were followed for 3 weeks post-inoculation. Mortality was recorded, and the lethal dose 50 (LD₅₀) was calculated as previously described [4].

RNA techniques. RNA was isolated as previously described [10], with slight modifications. Ten ml of a *S. suis* culture in the mid-exponential growth phase (OD₆₀₀ = 0.6) were collected by centrifugation at 8000 ×g for 10 min. Cells were then resuspended by vigorous shaking at 37°C for 10 min in 300 µl of Tris-EDTA pre-lysis buffer containing 10 mg of lysozyme per ml. Total RNA was extracted using the RNeasy Mini Kit (Qiagen), according to the manufacturer's instructions. Contaminating DNA was removed from the RNA during purification by treatment with RNase-free DNase (Qiagen) followed by digestion with DNase Turbo (Ambion). The concentration and integrity of RNA were determined by measuring the absorbance at 260 nm (A₂₆₀) and by 1% agarose gel electrophoresis, respectively. In all cases, the absence of DNA in RNA samples was confirmed by PCR without the addition of reverse transcriptase. Reverse transcription-PCR (RT-PCR)

Table 1. Bacterial strains and plasmids used in this work

Strain or plasmid	Relevant features	Source or reference
Strains		
<i>Streptococcus suis</i>		
P1/7	Virulent serotype 2 strain	
UA5001	Derived from P1/7. Δ <i>fur</i>	[3]
UA5003	Derived from P1/7. <i>feoB::spc</i> . <i>Spc</i> ^r	This work
<i>Escherichia coli</i>		
BL21-CodonPlus(DE3)-RIL	B ⁻ F ⁻ ompT hsdS (r _B m _B ⁻) dcm ⁺ Tc ^r gall (DE3) endA The [argU ileY leuW Cam ^r]	Stratagene
DH5α	<i>supE4 ΔlacU169 (Φ80 lacZΔM15) hsdR recA1 endA1 gyrA96 thi-1 relA1</i>	Clontech
Plasmids		
pGEM-T	PCR cloning vector. Ap ^r	Promega
pET15b	His ₆ tag expression vector. Ap ^r	Novagen
pR412	pEMcat derivative. Ap ^r , <i>Spc</i> ^r	[18]
pSET5s	Temperature-sensitive suicide vector for <i>S. suis</i> mutagenesis. Cat ^r	[28]
pSET5s <i>feoB::spc</i>	pSET5s vector carrying the construction for <i>feoB</i> mutagenesis	This work

Table 2. Oligonucleotide primers used in this work

Oligonucleotide	Sequence (5'–3')	Application
1638intUP	CTAGATTTGCCTGAATAAGG	Primer used to construct the pSET5 <i>feoB</i> :: <i>spc</i> plasmid
1638intDW	CTTGATGGCATTATTGCAGG	Primer used to construct the pSET5 <i>feoB</i> :: <i>spc</i> plasmid
1638upSma	CCCGGGATCTGCAAAACCTTCAAGTTGG	Primer used to construct the pSET5 <i>feoB</i> :: <i>spc</i> plasmid
1638dwSma	CCCGGGATTTCGGTAAATGCTAGTCTTG C	Primer used to construct the pSET5 <i>feoB</i> :: <i>spc</i> plasmid
SecAup2	TTCCAACGCCAAACTGG	Primer used to confirm <i>feoB</i> mutant
SecAup3	ACCGACTGAGCTAAACCAG	Primer used to confirm <i>feoB</i> mutant
M13F	GTTTTCCAGTCACGAC	Forward primer to amplify from pGEM-T vector. DIG-labelled at its 5' end to obtain EMSA probes
M13R	CAGGAAACAGCTATGAC	Reverse primer to amplify from pGEM-T vector. DIG-labelled at its 5' end to obtain EMSA probes
T7 Promoter	TAATACGACTCACTATAGGG	Forward primer for sequencing pET15b vector
T7 Terminator	GCTAGTTATTGCTCAGCGG	Reverse primer for sequencing pET15b vector
1637.up	ACACTTAGCGCACTTGGGTTTA	Upper primer for real-time quantitative RT-PCR assay to determine <i>feoA</i> gene expression and the upper primer used in RT-PCR assay
1637.dw	GCCTAGGGGAGCGACTTTTT	Lower primer for real-time quantitative RT-PCR assay to determine <i>feoA</i> gene expression
1638.up	CTGGCTGGCTACAATCTCTGGT	Upper primer for real-time quantitative RT-PCR assay to determine <i>feoB</i> gene expression
1638.dw	TTTTCGGTCCTGTTCAATTTCA	Lower primer for real-time quantitative RT-PCR assay to determine <i>feoB</i> gene expression and the lower primer used in RT-PCR assay
cys.up	GGTAACACTGGTATCGGTCTTG	Forward primer for real-time quantitative RT-PCR assay of the cysteine synthase gene
cys.dw	CCTGTCTGTGTATCTTCGTG	Reverse primer for real-time quantitative RT-PCR assay of the cysteine synthase gene
1637.pr	TCCTCTTAGGTGTTAGC	Forward primer to obtain EMSA probes of the <i>feoAB</i> promoter region
1637.50	TATCATCTTGCCTTAGGC	Reverse primer to obtain EMSA probes of the <i>feoAB</i> promoter region
fur.up.NdeI	ACTGACATATGGAAGTCCATTCTCACTT CAATGC	Upper primer for cloning the <i>S. suis fur</i> gene into the pET15b expression vector
fur.rv.BamHI	AGGGATCCATTCGTCTTTTAGCCCTGAC	Lower primer for cloning the <i>S. suis fur</i> gene into the pET15b expression vector

assays were done using a Titan One Tube RT-PCR system (Roche) and following the manufacturer's instructions. Real-time quantitative RT-PCR (rRT-PCR) analysis of total RNA was performed for all genes, as reported previously [6], using specific internal oligonucleotides for each gene (Table 2). The cysteine synthase gene was used as a standard, since its expression was not affected under the evaluated conditions. The total RNA concentrations of the WT and *fur* mutant strains were adjusted to the same value, and the genes to be tested, as well as the standard, were assayed simultaneously using a set of standard samples in each case.

Protein overexpression. The *S. suis* Fur protein was overexpressed as follows. The DNA fragment containing the *fur* gene was amplified from purified chromosomal DNA of *S. suis* P1/7 strain by PCR, using the primers fur.up.NdeI and fur.rv.BamHI (Table 2). The purified PCR product was then enzymatically digested with the corresponding restriction enzymes, cloned into the pET15b expression vector, and transformed into *E. coli* DH5 α cells. Recombinant plasmids were predicted to express an N-terminal His6-tagged fusion protein, and correct in-frame fusions of the protein into pET15b were confirmed by sequencing of plasmid DNA with T7 promoter and T7 termi-

nator primers (Macrogen Sequencing Service) (Table 2). Recombinant plasmids were then used to transform the BL21-CodonPlus (DE3)-RIL strain. An overnight culture of the resulting transformants was diluted (1/20) in 10 ml of LB medium and incubated at 37°C until OD₆₀₀ 0.6 was reached. Expression of the fusion protein was induced at this time by the addition of IPTG to a final concentration of 1 mM. After incubation for an additional 3 h at 37°C, the cells were collected by centrifugation at 8000 \times g for 10 min and resuspended in 20 mM Tris-HCl (pH 8), 50 mM KCl, 5% (vol/vol) glycerol, and a Complete Mini protease inhibitor cocktail (Roche). The cell suspension was then lysed by sonication on ice for 5 min at 50 W using a Braun LabsonicU (Braun Biotech). Unbroken cells and debris were removed by centrifugation at 7000 \times g for 10 min. Crude extracts were visualized by SDS-PAGE [16] and stored at 4°C for electrophoretic mobility shift assays .

Electrophoretic mobility shift assays (EMSA). EMSA were done as previously described [5] with slight modifications. The *feo* promoter was PCR-amplified from P1/7 *S. suis* genomic DNA using oligonucleotides 1637.pr and 1637.50 (Table 2). The obtained DNA fragment (396 bp), extending from position +67 to –329 (with respect to its predicted transla-




Fig. 1. Genetic organization of *Streptococcus suis* P1/7 strain chromosomal region containing the *feoAB* genes, based on [http://www.sanger.ac.uk/Projects/S_suis/]. The large black arrow indicates the putative transcriptional unit that was tested. The primer sets used for the RT-PCR assays are indicated by small arrows. RT-PCR was carried out in the presence of total RNA (lane A), DNA (lane B), or in the absence of both RNA and DNA (lane C). *Bst*EII-digested- λ DNA was used as a molecular size marker (MW). The lengths of the gene products in number of amino acids and the putative functions of the genes are also indicated.

tional starting point), was cloned in *E. coli* DH5 α cells through the pGEM-T vector. The presence of the desired promoter was confirmed by sequencing the plasmid DNA using the primers M13F and M13R (Table 2). DNA probes were prepared by PCR amplification, with one of the oligonucleotides labeled with digoxigenin (DIG) at its 5' end. DNA-protein reaction mixtures (20 μ l) containing 25 ng of a DIG-DNA-labeled probe and different concentrations of either crude extracts overexpressing or not the *S. suis* Fur protein were incubated in EMSA buffer [20 mM Tris-HCl (pH 8), 50 mM KCl, 5% (vol/vol) glycerol, 1 μ g bulk carrier sperm salmon DNA, 0.5 mM 1,4-dithiothreitol, and 0.1 mg bovine serum albumin per ml] for 10 min at room temperature. DNA-protein complexes were visualized by separation on a 5% nondenaturing polyacrylamide gel (40 mM Tris-acetate) at 150 V for 1.5 h, and then transferred to a Biodine B nylon membrane (Pall Gelman Laboratory). DIG-DNA-labeled-protein complexes were detected according to the manufacturer's protocol (Roche).


Results and Discussion

The upstream region of the *E. coli* *feo* operon contains a binding site for the Fur protein. Together with iron (Fe²⁺), Fur acts as a corepressor [15]. In a previous work, the *E. coli* (*fur*) homologue was identified in the *S. suis* P1/7 strain through bioinformatics analysis [3]. In this study, further *in silico* searches in the database containing the complete annotation of the *S. suis* P1/7 strain genome [http://www.sanger.ac.uk/Projects/S_suis/] were carried out using the TBLASTX program and the *E. coli* *feoA* gene sequence as probe.

An open reading frame (ORF) corresponding to a protein of 156 amino acid residues and homologous to the *E. coli* *feoA* gene was found to be encoded by the *S. suis* P1/7 strain SSU_1236 gene. The calculated molecular mass of the predicted translated *S. suis* FeoA protein was 17,433 Da. Immediately downstream of the SSU_1236 gene, another ORF (encoded by gene SSU_1237), homologous to the *E. coli* *feoB* gene, was identified (Fig. 1). The predicted translated FeoB protein (714 amino acid residues) had a calculated molecular mass of 79,083 Da. In contrast to the organization of the gene in *E. coli*, no *feoC* homologues were found in the *S. suis* genome. The identity between the *S. suis* FeoA and FeoB proteins and the protein products encoded by the *E. coli* *feoA* and *feoB* genes was 37% and 35%, respectively. RT-PCR analysis of the transcriptional pattern of the *S. suis* *feoAB* genes showed that they belong to the same transcriptional unit (Fig. 1). A similar transcriptional organization of the *feo* genes is found in *L. monocytogenes* [14,15].

In silico analysis of the upstream regions of *feo* operons from multiple proteobacterial genomes have revealed a conserved palindromic motif corresponding to a Fur binding site [9,23]. However, no apparent conserved motifs were found in the promoter region of the *feoAB* *S. suis* operon. Since in *E. coli* the *feoABC* operon is under the control of Fur [15], we overexpressed the *S. suis* Fur protein (Fig. 2A) and then used

Fig. 2. (A) SDS-PAGE profile of a crude extract from *Escherichia coli* BL21-CodonPlus(DE3)-RIL cells containing the pET15b vector without (lane 1) or with (lane 2) overexpression of the *Streptococcus suis* Fur 17-kDa protein. (B) Electrophoretic mobility of DNA fragments containing the *S. suis* *feoAB* promoter. The analyses were carried out in the presence of increasing amounts (0–10 µg) of crude extract from *E. coli* BL21-CodonPlus(DE3)-RIL cells containing (left) or not (right) the pET15b vector overexpressing the *S. suis* Fur protein. All EMSA were repeated a minimum of three times to ensure the reproducibility of the results. (C) Up-regulation of *feoA* and *feoB* genes in the *S. suis* *fur* mutant (UA5001) as measured by real-time quantitative RT-PCR. The expression factor is the ratio of the mRNA concentration of each gene from UA5001 strain with respect to the WT parent strain P1/7 (defined as 1). The relative mRNA concentration of each gene was determined by using a standard curve generated by amplifying an internal fragment of the *S. suis* P1/7 ORF encoding the cysteine synthase enzyme. The mean value and the standard deviation from two independent experiments (each in duplicate) are also indicated.



EMSA studies to test its ability to bind the promoter region of the *feoAB* transcriptional unit. Interestingly, the EMSA results showed that *S. suis* Fur binds to the *feoAB* promoter (Fig. 2B). Binding was specific since the presence of unrelated DNA as a competitor did not abolish the Fur-mediated retarding band (Fig. 2B). In agreement with these data, derepression of *feoA* and *feoB* gene expression was observed in the *S. suis* *fur* mutant UA5001 [3] through real-time quantitative RT-PCR assays (Fig. 2C).

In order to study the contribution of the *feoAB* operon to *S. suis* virulence, the *S. suis* *feoB* gene was inactivated in the P1/7 virulent strain, giving rise to mutant strain UA5003 *feoB::spc* (Table 1). The virulence of this strain was subsequently evaluated in a murine model of infection, comparing the LD₅₀ of the *feoB::spc* mutant to that of the WT parent strain P1/7. Four groups of three female 8-week-old BALB/cAnNHsd mice were injected intraperitoneally with 0.1 ml of serial 10-fold dilutions of bacteria prepared from each bacterial strain. This assay showed that the virulence of the *feoB* mutant (LD₅₀ = 3.9 × 10⁷ CFU/animal) was signifi-

cantly attenuated compared with that of the WT strain (LD₅₀ = 4.4 × 10⁶ CFU/animal). Note that the in vitro growth rates of the *S. suis* *feoB* mutant and the WT strains did not differ under the metal-repleted conditions present in the TH medium (Fig. 3). However, in the presence of the chelator DPD, the growth rate of the *feoB* mutant was lower than that of the wild-type P1/7 strain. This behavior can be attributed to a decreased capacity of the FeoB-deficient mutant to grow under metal-depleted conditions.

Streptococcus suis has been shown to not require iron for growth in vitro [21]. In addition, the degree of identity between FeoB from *S. suis* and the *E. coli* Feo transporter is low, and some FeoB homologues transport manganese rather than iron [8]. However, the impaired virulence of the *S. suis* *feoB* mutant, together with the fact that the genes encoding the Feo transporter are Fur regulated, indicate that, under the low concentrations of these elements existing inside host cells and in mammalian body fluids, the *feoB*-encoded protein of *S. suis* might act as an important transporter for metal uptake in vivo. Furthermore, it has been reported that the




Fig. 3. Growth of wild-type P1/7 (open squares) and UA5003 *feoB* mutant (closed squares) of *Streptococcus suis* strains in TH medium; and the same strains, P1/7 (closed triangles) and UA5003 (open triangles), in TH medium in the presence of the metal chelator DPD (100 μ M). The means from three independent experiments (for each growth curve) are represented. For simplification, error bars have been omitted; the standard deviation was <10 % in all cases.

FeoB transporter is required for the intracellular growth of *Legionella pneumophila* [22] and for the virulence of several gram-negative bacteria [8,20,22,25,29].

To summarize, the data reported in this work demonstrate that full virulence of *S. suis* requires the FeoB transporter. To our knowledge, this study is the first to describe the contribution of the *feoB* gene to the virulence of gram-positive bacterial pathogens.

Acknowledgements. We are grateful to Dr. Sekizaki for providing plasmid pSET5s. This work was funded by grants AGL2005-03574, from the Ministerio de Educación y Ciencia, and 2005SGR-533, from DURSI of the Generalitat de Catalunya. N.F. has a Natural Sciences and Engineering Research Council of Canada postgraduate scholarship.

References

- Andrews SC, Robinson AK, Rodriguez-Quinones F (2003) Bacterial iron homeostasis. *FEMS Microbiol Rev* 27:215-237
- Aranda J, Garrido ME, Cortés P, Llagostera M, Barbé J (2008) Analysis of the protective capacity of three *Streptococcus suis* proteins induced under divalent-cation-limited conditions. *Infect Immun* 76:1590-1598
- Aranda J, Garrido ME, Fittipaldi N, Cortes P, Llagostera M, Gottschalk M, Barbe J (2009) Protective capacities of cell surface-associated proteins of *Streptococcus suis* mutants deficient in divalent-cation-uptake regulators. *Microbiology* 155:1580-1587
- Bigas A, Garrido ME, Badiola I, Barbe J, Llagostera M (2006) Colonization capacity and serum bactericidal activity of *Haemophilus parasuis* thy mutants. *Int Microbiol* 9:297-301
- Bsat N, Helmann JD (1999) Interaction of *Bacillus subtilis* Fur (ferric uptake repressor) with the *dhb* operator in vitro and in vivo. *J Bacteriol* 181:4299-4307
- Campoy S, Mazón G, Fernández de Henestrosa AR, Llagostera M, Monteiro PB, Barbé J (2002) A new regulatory DNA motif of the gamma subclass Proteobacteria: identification of the LexA protein binding site of the plant pathogen *Xylella fastidiosa*. *Microbiology* 148:3583-3597
- Cartron ML, Maddocks S, Gillingham P, Craven CJ, Andrews SC (2006) Feo-transport of ferrous iron into bacteria. *Biometals* 19:143-157
- Dashper SG, Butler CA, Lissel JP, et al. (2005) A novel *Porphyrromonas gingivalis* FeoB plays a role in manganese accumulation. *J Biol Chem* 280:28095-28102
- Escolar L, Pérez-Martín J, de Lorenzo V (1999) Opening the iron box: transcriptional metalloregulation by the Fur protein. *J Bacteriol* 181:6223-6229
- Fontaine MC, Perez-Casal J, Willson PJ (2004) Investigation of a novel DNase of *Streptococcus suis* serotype 2. *Infect Immun* 72:774-781
- Garrido ME, Bosch M, Bigas A, Badiola I, Barbé J, Llagostera M (2008) Heterologous protective immunization elicited in mice by *Pasteurella multocida fur ompH*. *Int Microbiol* 11:17-24
- Gottschalk M, Segura M, Xu J (2007) *Streptococcus suis* infections in humans: the Chinese experience and the situation in North America. *Anim Health Res Rev* 8:29-45
- Higgins R, Gottschalk M (2005) Streptococcal diseases. In: Straw BE, et al. (eds) *Diseases of swine*. Iowa State University Press, Ames, Iowa
- Jin B, Newton SM, Shao Y, Jiang X, Charbit A, Klebba PE (2006) Iron acquisition systems for ferric hydroxamates, haemin and haemoglobin in *Listeria monocytogenes*. *Mol Microbiol* 59:1185-1198

15. Kammler M, Schon C, Hantke K (1993) Characterization of the ferrous iron uptake system of *Escherichia coli*. *J Bacteriol* 175:6212-6219
16. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685
17. Mai NT, Hoa NT, Nga TV, et al. (2008) *Streptococcus suis* meningitis in adults in Vietnam. *Clin Infect Dis*. 46:659-667
18. Martin B, Prudhomme M, Alloing G, Granadel C, Claverys JP (2000) Cross-regulation of competence pheromone production and export in the early control of transformation in *Streptococcus pneumoniae*. *Mol Microbiol* 38:867-878
19. Miller JH (1992) A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
20. Naikare H, Palyada K, Panciera R, Marlow D, Stintzi A (2006) Major role for FeoB in *Campylobacter jejuni* ferrous iron acquisition, gut colonization, and intracellular survival. *Infect Immun* 74:5433-5444
21. Niven DF, Ekins A, al-Samaurai AA (1999) Effects of iron and manganese availability on growth and production of superoxide dismutase by *Streptococcus suis*. *Can J Microbiol* 45:1027-1032
22. Robey M, Cianciotto NP (2002) *Legionella pneumophila feoAB* promotes ferrous iron uptake and intracellular infection. *Infect Immun* 70:5659-5669
23. Rodionov DA, Dubchak I, Arkin A, Alm E, Gelfand MS (2004) Reconstruction of regulatory and metabolic pathways in metal-reducing δ -proteobacteria. *Genome Biol* 5:R90
24. Sambrook J, Russell DW (2001) Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
25. Stojiljkovic I, Cobeljic M, Hantke K (1993) *Escherichia coli* K-12 ferrous iron uptake mutants are impaired in their ability to colonize the mouse intestine. *FEMS Microbiol Lett* 108:111-115
26. Suankratay C, Intalaporn P, Nunthapisud P, Arunyingmongkol K, Wilde H (2004) *Streptococcus suis* meningitis in Thailand. *Southeast Asian J Trop Med Public Health* 35:868-876
27. Takamatsu D, Osaki M, Sekizaki T (2001) Construction and characterization of *Streptococcus suis*-*Escherichia coli* shuttle cloning vectors. *Plasmid* 45:101-113
28. Takamatsu D, Osaki M, Sekizaki T (2001) Thermosensitive suicide vectors for gene replacement in *Streptococcus suis*. *Plasmid* 46:140-148
29. Velayudhan J, Hughes NJ, McColm AA, Bagshaw J, Clayton CL, Andrews SC, Kelly DJ (2000) Iron acquisition and virulence in *Helicobacter pylori*: a major role for FeoB, a high-affinity ferrous iron transporter. *Mol Microbiol* 37:274-286
30. Wandersman C, Delepelaire P (2004) Bacterial iron sources: from siderophores to hemophores. *Annu Rev Microbiol* 58:611-647